



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Completing squares, $x^4 + 420x^2 + 210^2 = 441x^2 + 8820x + 210^2$, or $(x^2 + 210)^2 = (21x + 210)^2$.

Whence, $x^2 = 21x$, $x = 0$, the introduced root, 21, and $\frac{1}{2}[-21 \pm \sqrt{-1239}]$. Hence, 21 is the only possible real value for x . Therefore, $2940/21 = 140$, the other factor. Hence 21 and 140 are the factors.

Also solved by H. Prime, V. M. Spunar, S. Lefschetz, J. Scheffer, and S. G. Barton.

GEOMETRY.

377. Proposed by S. A. COREY, Hiteman, Iowa.

Let AB, BC, CD, DE, EA be the sides of a pentagon, plane or gauche. From A draw AF, AG, AH , parallel to, and of the same length and currency as BC, CD, DE , respectively. Bisect AE at K . Draw KB, KF, KG , and KH . Prove that $KB^2 + KF^2 + KG^2 + KH^2 = AB^2 + BC^2 + CD^2 + DE^2$.

I. Solution by J. SCHEFFER, A. M., Hagerstown, Md.

K being the mid-point of the diagonal AE in the parallelogram $ADEH$, DKA must be a straight line, viz., the other diagonal. Let the orthogonal coordinates of B, F, G, H , with reference to AE as the axis of X , and A as origin, be, respectively, $x_1, y_1; x_2, y_2; x_3, y_3; x_4, y_4$; and $AK = EK = a$. Then


$$AB^2 + AF^2 + AG^2 + AH^2 = (x_1^2 + y_1^2) + (x_2^2 + y_2^2) + (x_3^2 + y_3^2) + (x_4^2 + y_4^2) \dots (I),$$

$$\text{and } KB^2 + KF^2 + KG^2 + KH^2$$

$$\begin{aligned} &= [(x_1 - a)^2 + y_1^2] + [(x_2 - a)^2 + y_2^2] + [(x_3 - a)^2 + y_3^2] + [(x_4 - a)^2 + y_4^2] \\ &= (x_1^2 + y_1^2) + (x_2^2 + y_2^2) + (x_3^2 + y_3^2) + (x_4^2 + y_4^2) \\ &\quad - 2a[(x_1 + x_2 + x_3 + x_4) - 2a] \dots (II). \end{aligned}$$

Letting fall the perpendiculars BM, CN, DO upon AE , and BP, DQ upon CN , we have $AB \cos BAM + BC \cos CBP + CD \cos CDA + DE \cos DEO = AB \cos BAM + AF \cos FAK + AG \cos KAG + AH \cos KAH = AE = 2a$.

$\therefore x_1 + x_2 + x_3 + x_4 = 2a$; therefore in (II), $x_1 + x_2 + x_3 + x_4 - 2a = 0$, and $AB^2 + AF^2 + AG^2 + AH^2 = KB^2 + KF^2 + KG^2 + KH^2 = AB^2 + BC^2 + CD^2 + DE^2$. Q. E. D.



II. Solution by the PROPOSER.

Let a , b , c , and d be the vector sides AB , BC , CD , and DE , respectively. Then will $EA = -(a+b+c+d)$, $AK = \frac{1}{2}(a+b+c+d)$, $KB = \frac{1}{2}(a-b-c-d)$, $KF = \frac{1}{2}(-a+b-c-d)$, $KG = \frac{1}{2}(-a-b+c-d)$, and $KH = \frac{1}{2}(-a-b-c+d)$.

Squaring these vector expressions for KB , KF , KG , and KH , and adding, their sum is found to be $a^2 + b^2 + c^2 + d^2$. As the square of a vector equals minus the square of its tensor, the truth of the proposition is demonstrated. Observe that a line drawn from D to K is equal to and may be substituted for KH in the equation of the problem.

378. Proposed by G. I. HOPKINS, A. M., Instructor in Mathematics and Astronomy, Manchester High School Manchester, N. H.

In the triangle AED , the lines BE and CE are drawn to the points B and C in the base of the triangle. If $AE=100$, $ED=125$, $BC=60$, and $\angle AEC = \angle BED =$ a right angle, compute AB , BE , EC , and CD .

Solution by A. H. HOLMES, Brunswick, Maine.

Put $DAE = \theta$ and $ADE = \psi$. Let fall perpendicular EF on base AD . Then we have, $100\sin \theta = 125\sin \psi$ or $4\sin \theta = 5\sin \psi \dots (1)$.

Since BED and AEC are right angles, $BD = \frac{DE}{\cos \psi}$, and $AC = \frac{AE}{\cos \theta}$.

$$\therefore \frac{100}{\cos \theta} + \frac{125}{\cos \psi} - 60 = 100\cos \theta + 125\cos \psi \dots (2).$$

Eliminating $\sin \theta$ and $\cos \psi$ from (1) and (2), and reducing,
 $\cos^6 \psi + .96\cos^5 \psi - 3.1296\cos^4 \psi - .0256\cos^3 \psi + 1.94745\cos^2 \psi - .1152\cos \psi = 0$.

Solving by Horner's method, $\cos \psi = .8851 +$. $\therefore \cos \theta = .8134 +$.

Then since $AC = 122.94 +$, $AB = 62.94 +$.

Similarly, $CD = 81.22 +$. Also, $BE = 65.71 +$, and $CE = 71.51 +$.

Also solved by J. Scheffer.

382. Proposed by PROF. R. C. ARCHIBALD, Brown University, Providence, R. I.

Between the side of a given rhombus and its adjacent side produced, to insert a straight line of a given length and directed to the opposite corner. "Euclidean constructions" are particularly desired.

Remark by V. M. SPUNAR, M. and E. E., Chicago, Illinois.

This is the famous Pappus problem: Rhombo dato et uno latere producto aptare sub angelo exteriori magnitudine datum rectam lineam, quae ad oppositum angulum pertingat.

Pappus, and a certain number of mathematicians, among them Newton, Huygens, and Gergonne, solved the problem algebraically and geometrically. (See E. Pruvost, *Geométrie Analytique*, t. I, pp. 18-28.)

The problem in the present form, proposed by Prof. R. C. Archibald